arXiv:1401.3161v1 [math.NT] 14 Jan 2014

IDENTITIES OF SYMMETRY FOR EXPANSIONS OF q-EULER POLYNOMIALS

DAE SAN KIM AND TAE GYUN KIM

Abstract. Recently, Kim considered expansions of q-Euler polynomials which are given by

$$\sum_{n=0}^{\infty} E_{n,q}^{(h,r)}(x)\frac{t^n}{n!} = 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} q^{\sum_{l=1}^{r}(h-l)m_l}(-1)^{\sum_{l=1}^{r} m_l} e^{[x+\sum_{l=1}^{r} m_l]_q t}.$$

In this paper, we investigate some symmetric properties of the multivariate p-adic fermionic integrals on $\mathbb{Z}_p$ and derive various identities concerning the expansions of q-Euler polynomials from the symmetric properties of the multivariate p-adic fermionic integrals on $\mathbb{Z}_p$

1. Introduction

Let p be a fixed odd prime number. Throughout this paper $\mathbb{Z}_p, \mathbb{Q}_p, \mathbb{C}_p$ will, respectively, denote the ring of p-adic rational integers, the field of p-adic rational numbers and the completion of algebraic closure of $\mathbb{Q}_p$. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = p^{-1}$. When one talks about a q-extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$ or a p-adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one usually assumes $|q| < 1$;if $q \in \mathbb{C}_p$, one usually assumes $|1-q|_p < 1$. The q-number of x is defined by $[x]_q = \frac{1-q^x}{1-q}$. Note that $\lim_{q\to 1}[x]_q = x$. The q-factorial is defined as $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$. As is well known, the q-binomial formulae are given by

$$\begin{aligned}(b;q)_n &= (1-b)(1-bq)\cdots(1-bq^{n-1}) \\ &= \sum_{i=0}^{n} \binom{n}{i}_q q^{\binom{i}{2}}(-1)^i b^i, \text{ (see [1, 5])},\end{aligned} \tag{1.1}$$

where $\binom{n}{i}_q = \frac{[n]_q!}{[i]_q![n-i]_q!} = \frac{[n]_q\cdots[n-i+1]_q}{[i]_q!}$, and

$$\frac{1}{(b;q)_n} = \frac{1}{(1-b)(1-bq)\cdots(1-bq^{n-1})} = \sum_{i=0}^{\infty} \binom{n+i-1}{i}_q b^i. \tag{1.2}$$

Let $C(\mathbb{Z}_p)$ be the space of continuous functions on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, the p-adic fermionic integral on $\mathbb{Z}_p$ is defined by Kim as follows:

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x)d\mu_{-1}(x) = \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)(-1)^x, \text{ (see [6])}. \tag{1.3}$$

From (1.3), we note that

$$I_{-1}(f_n)+(-1)^{n-1}I_{-1}(f)=2\sum_{l=0}^{n-1}(-1)^{n-l-1}f(l), \tag{1.4}$$

where $f_n(x)=f(x+n)$ and $n\in\mathbb{N}$.
As is well known, the higher-order Euler polynomials are defined by the generating function to be

$$\left(\frac{2}{e^t+1}\right)^r e^{xt}=\sum_{n=0}^{\infty}E_n^{(r)}(x)\frac{t^n}{n!},\ (r\in\mathbb{N}),\ \text{(see [1-14])}. \tag{1.5}$$

The q-extension of (1.5) is given by

$$\begin{aligned}\sum_{n=0}^{\infty}E_{n,q}^{(r)}(x)\frac{t^n}{n!}&=2^r\sum_{m_1,\cdots,m_r=0}^{\infty}(-1)^{m_1+\cdots+m_r}e^{[m_1+\cdots+m_r+x]_q t}\\&=2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}(-1)^m e^{[m+x]_q t},\ \text{(see [5,6])}.\end{aligned} \tag{1.6}$$

Recently, Kim considered expansions of $E_{n,q}^{(r)}(x)$ which are given by

$$\begin{aligned}\sum_{n=0}^{\infty}E_{n,q}^{(h,r)}(x)\frac{t^n}{n!}&=2^r\sum_{m_1,\cdots,m_r=0}^{\infty}q^{\sum_{l=1}^{r}(h-l)m_l}(-1)^{\sum_{l=1}^{r}m_l}e^{[x+m_1+\cdots+m_r]_q t}\\&=2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}_q q^{(h-r)m}(-1)^m e^{[m+x]_q t},\end{aligned} \tag{1.7}$$

where $h\in\mathbb{Z}$ and $r\in\mathbb{N}$ (see [5,6,7]).
In this paper, we investigate some symmetric properties of the multivariate p-adic fermionic integrals on $\mathbb{Z}_p$ and derive various identities concerning the expansions of q-Euler polynomials from the symmetric properties of the multivariate p-adic fermionic interals on $\mathbb{Z}_p$.

2. Identities of Symmetry for expansions of q-Euler polynomials

For $h\in\mathbb{Z}$ and $r\in\mathbb{N}$, from (1.3), we have

$$\begin{aligned}&\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{\sum_{l=1}^{r}(h-l)y_l}e^{[x+y_1+\cdots+y_r]_q t}d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)\\&=2^r\sum_{m_1,\cdots,m_r=0}^{\infty}q^{\sum_{l=1}^{r}(h-l)m_l}(-1)^{\sum_{l=1}^{r}m_l}e^{[x+m_1+\cdots+m_r]_q t}\\&=2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}_q q^{(h-r)m}(-1)^m e^{[m+x]_q t}.\end{aligned} \tag{2.1}$$

Thus, by (1.7) and (2.1), we get

$$\begin{aligned}
E_{n,q}^{(h,r)}(x) &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{\sum_{l=1}^{r}(h-l)y_l}[x+y_1+\cdots+y_r]_q^n d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r) \\
&= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-q^{h-r})^m [x+m]_q^n \\
&= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \frac{\binom{n}{l}(-q^x)^l}{(-q^{h-r+l};q)_r} \\
&= 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} q^{\sum_{l=1}^{r}(h-l)m_l}(-1)^{\sum_{l=1}^{r} m_l}[m_1+\cdots+m_r+x]_q^n,
\end{aligned} \tag{2.2}$$

where $n \geq 0$.

Let $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$ and $w_2 \equiv 1 \pmod 2$. Then we see that

$$\begin{aligned}
&\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{w_1\sum_{l=1}^{r}(h-l)y_l} e^{[w_1]_q[w_2x+\frac{w_2}{w_1}\sum_{l=1}^{r} j_l+\sum_{l=1}^{r} y_l]_{q^{w_1}}t} d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r) \\
&= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{w_1\sum_{l=1}^{r}(h-l)y_l} e^{[w_1w_2x+w_2\sum_{l=1}^{r} j_l+w_1\sum_{l=1}^{r} y_l]_q t} d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r) \\
&= \lim_{N\to\infty} \sum_{y_1,\cdots,y_r=0}^{w_2p^N-1} q^{w_1\sum_{l=1}^{r}(h-l)y_l} e^{[w_1w_2x+w_2\sum_{l=1}^{r} j_l+w_1\sum_{l=1}^{r} y_l]_q t}(-1)^{\sum_{l=1}^{r} y_l} \\
&\qquad\qquad \times d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r) \\
&= \lim_{N\to\infty} \sum_{i_1,\cdots,i_r=0}^{w_2-1} \sum_{y_1,\cdots,y_r=0}^{p^N-1} q^{w_1\sum_{l=1}^{r}(h-l)(i_l+w_2y_l)} \\
&\qquad\qquad \times e^{[w_1w_2x+w_2\sum_{l=1}^{r} j_l+w_1\sum_{l=1}^{r}(i_l+w_2y_l)]_q t}(-1)^{\sum_{l=1}^{r}(i_l+w_2y_l)}.
\end{aligned} \tag{2.3}$$

From (2.3), we note that

$$\begin{aligned}
&\sum_{j_1,\cdots,j_r=0}^{w_1-1} (-1)^{\sum_{l=1}^{r} j_l} q^{w_2\sum_{l=1}^{r}(h-l)j_l} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{w_1\sum_{l=1}^{r}(h-l)y_l} \\
&\qquad\qquad \times e^{[w_1]_q[w_2x+\frac{w_2}{w_1}\sum_{l=1}^{r} j_l+\sum_{l=1}^{r} y_l]_{q^{w_1}}t} d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r) \\
&= \lim_{N\to\infty} \sum_{j_1,\cdots,j_r=0}^{w_1-1} \sum_{i_1,\cdots,i_r=0}^{w_2-1} \sum_{y_1,\cdots,y_r=0}^{p^N-1} (-1)^{\sum_{l=1}^{r}(i_l+j_l+y_l)} q^{\sum_{l=1}^{r}(h-l)(w_2j_l+w_1i_l+w_1w_2y_l)} \\
&\qquad\qquad \times e^{[w_1w_2(x+\sum_{l=1}^{r} y_l)+\sum_{l=1}^{r} w_2j_l+w_1i_l]_q t}.
\end{aligned} \tag{2.4}$$

By the same method as (2.4), we get

$$\sum_{j_1,\cdots,j_r=0}^{w_2-1}(-1)^{\sum_{l=1}^r j_l}q^{w_1\sum_{l=1}^r(h-l)j_l}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{w_2\sum_{l=1}^r(h-l)y_l}$$
$$\times e^{[w_2]_q[w_1x+\frac{w_1}{w_2}\sum_{l=1}^r j_l+\sum_{l=1}^r y_l]_{q^{w_2}}t}d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)$$
$$=\lim_{N\to\infty}\sum_{j_1,\cdots,j_r=0}^{w_2-1}\sum_{i_1,\cdots,i_r=0}^{w_1-1}\sum_{y_1,\cdots,y_r=0}^{p^N-1}(-1)^{\sum_{l=1}^r(i_l+j_l+y_l)}q^{\sum_{l=1}^r(h-l)(w_1j_l+w_2i_l+w_1w_2y_l)}$$
$$\times e^{[w_1w_2(x+\sum_{l=1}^r y_l)+\sum_{l=1}^r w_1j_l+w_2i_l]_q t} \tag{2.5}$$

Therefore, by (2.4) and (2.5), we obtain the following theorem.

Theorem 2.1. *For $w_1,w_2\in\mathbb{N}$ with $w_1\equiv 1 \pmod 2$ and $w_2\equiv 1 \pmod 2$, we have*

$$\sum_{j_1,\cdots,j_r=0}^{w_1-1}(-1)^{\sum_{l=1}^r j_l}q^{w_2\sum_{l=1}^r(h-l)j_l}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{w_1\sum_{l=1}^r(h-l)y_l}$$
$$\times e^{[w_1]_q[w_2x+\frac{w_2}{w_1}\sum_{l=1}^r j_l+\sum_{l=1}^r y_l]_{q^{w_1}}t}d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)$$
$$=\sum_{j_1,\cdots,j_r=0}^{w_2-1}(-1)^{\sum_{l=1}^r j_l}q^{w_1\sum_{l=1}^r(h-l)j_l}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{w_2\sum_{l=1}^r(h-l)y_l}$$
$$\times e^{[w_2]_q[w_1x+\frac{w_1}{w_2}\sum_{l=1}^r j_l+\sum_{l=1}^r y_l]_{q^{w_2}}t}d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)$$

Corollary 2.2. *For $n\geq 0$, $w_1,w_2\in\mathbb{N}$ with $w_1\equiv 1 \pmod 2$ and $w_2\equiv 1 \pmod 2$, we have*

$$[w_1]_q^n\sum_{j_1,\cdots,j_r=0}^{w_1-1}(-1)^{\sum_{l=1}^r j_l}q^{w_2\sum_{l=1}^r(h-l)j_l}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{w_1\sum_{l=1}^r(h-l)y_l}$$
$$\times[w_2x+\frac{w_2}{w_1}\sum_{l=1}^r j_l+\sum_{l=1}^r y_l]_{q^{w_1}}^n d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)$$
$$=[w_2]_q^n\sum_{j_1,\cdots,j_r=0}^{w_2-1}(-1)^{\sum_{l=1}^r j_l}q^{w_1\sum_{l=1}^r(h-l)j_l}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{w_2\sum_{l=1}^r(h-l)y_l}$$
$$\times[w_1x+\frac{w_1}{w_2}\sum_{l=1}^r j_l+\sum_{l=1}^r y_l]_{q^{w_2}}^n d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)$$

Therefore, by (2.2) and Corollary 2.2, we obtain the following theorem.

Theorem 2.3. *For $n\geq 0$, $w_1,w_2\in\mathbb{N}$ with $w_1\equiv 1 \pmod 2$ and $w_2\equiv 1 \pmod 2$, we have*

$$[w_1]_q^n\sum_{j_1,\cdots,j_r=0}^{w_1-1}(-1)^{\sum_{l=1}^r j_l}q^{w_2\sum_{l=1}^r(h-l)j_l}E_{n,q^{w_1}}^{(h,r)}(w_2x+\frac{w_2}{w_1}(j_1+\cdots+j_r))$$
$$=[w_2]_q^n\sum_{j_1,\cdots,j_r=0}^{w_2-1}(-1)^{\sum_{l=1}^r j_l}q^{w_1\sum_{l=1}^r(h-l)j_l}E_{n,q^{w_2}}^{(h,r)}(w_1x+\frac{w_1}{w_2}(j_1+\cdots+j_r)).$$

From (1.3), we note that

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p} q^{w_1\sum_{l=1}^{r}(h-l)y_l}[w_2x+\frac{w_2}{w_1}\sum_{l=1}^{r}j_l+\sum_{l=1}^{r}y_l]_{q^{w_1}}^{n}\,d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)\\
&=\sum_{i=0}^{n}\binom{n}{i}\left(\frac{[w_2]_q}{[w_1]_q}\right)^{i}[j_1+\cdots+j_r]_{q^{w_2}}^{i}q^{w_2(n-i)\sum_{l=1}^{r}j_l}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}[w_2x+\sum_{l=1}^{r}y_l]_{q^{w_1}}^{n-i}\\
&\qquad\qquad\times q^{w_1\sum_{l=1}^{r}(h-l)y_l}d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)\\
&=\sum_{i=0}^{n}\binom{n}{i}\left(\frac{[w_2]_q}{[w_1]_q}\right)^{i}[j_1+\cdots+j_r]_{q^{w_2}}^{i}q^{w_2(n-i)\sum_{l=1}^{r}j_l}E_{n-i,q^{w_1}}^{(h,r)}(w_2x).
\end{aligned}\tag{2.6}$$

By (2.6), we get

$$\begin{aligned}
&[w_1]_q^n\sum_{j_1,\cdots,j_r=0}^{w_1-1}(-1)^{\sum_{l=1}^{r}j_l}q^{w_2\sum_{l=1}^{r}(h-l)j_l}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{w_1\sum_{l=1}^{r}(h-l)y_l}\\
&\qquad\qquad\times[w_2x+\frac{w_2}{w_1}\sum_{l=1}^{r}j_l+\sum_{l=1}^{r}y_l]_{q^{w_1}}^{n}\,d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r)\\
&=\sum_{j_1,\cdots,j_r=0}^{w_1-1}(-1)^{\sum_{l=1}^{r}j_l}q^{w_2\sum_{l=1}^{r}(h-l)j_l}\sum_{i=0}^{n}\binom{n}{i}[w_2]_q^i[w_1]_q^{n-i}[j_1+\cdots+j_r]_{q^{w_2}}^{i}\\
&\qquad\qquad\times q^{w_2(n-1)\sum_{l=1}^{r}j_l}E_{n-i,q^{w_1}}^{(h,r)}(w_2x)\\
&=\sum_{i=0}^{n}\binom{n}{i}[w_2]_q^i[w_1]_q^{n-i}E_{n-i,q^{w_1}}^{(h,r)}(w_2x)\sum_{j_1,\cdots,j_r=0}^{w_1-1}(-1)^{\sum_{l=1}^{r}j_l}q^{w_2\sum_{l=1}^{r}(n+h-l-i)j_l}\\
&\qquad\qquad\times[j_1+\cdots+j_r]_{q^{w_2}}^{i}\\
&=\sum_{i=0}^{n}\binom{n}{i}[w_2]_q^i[w_1]_q^{n-i}E_{n-i,q^{w_1}}^{(h,r)}(w_2x)T_{n,i,q^{w_2}}^{(h,r)}(w_1),
\end{aligned}\tag{2.7}$$

where

$$T_{n,i,q}^{(h,r)}(w)=\sum_{j_1,\cdots,j_r=0}^{w-1}(-1)^{\sum_{l=1}^{r}j_l}q^{\sum_{l=1}^{r}(n+h-l-i)j_l}[j_1+\cdots+j_r]_q^i.\tag{2.8}$$

By the same method as (2.7), we get

$$
\begin{aligned}
&[w_2]_q^n \sum_{j_1,\cdots,j_r=0}^{w_2-1} (-1)^{\sum_{l=1}^r j_l} q^{w_1\sum_{l=1}^r (h-l)j_l} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{w_2\sum_{l=1}^r (h-l)y_l} \\
&\qquad \times [w_1 x + \frac{w_1}{w_2}\sum_{l=1}^r j_l + \sum_{l=1}^r y_l]_{q^{w_2}}^n d\mu_{-1}(y_1)\cdots d\mu_{-1}(y_r) \\
&= \sum_{j_1,\cdots,j_r=0}^{w_2-1} (-1)^{\sum_{l=1}^r j_l} q^{w_1\sum_{l=1}^r (h-l)j_l} \sum_{i=0}^n \binom{n}{i} [w_1]_q^i [w_2]_q^{n-i} [j_1+\cdots+j_r]_{q^{w_1}}^i \\
&\qquad \times q^{w_1(n-i)\sum_{l=1}^r j_l} E_{n-i,q^{w_2}}^{(h,r)}(w_1 x) \\
&= \sum_{i=0}^n \binom{n}{i} [w_1]_q^i [w_2]_q^{n-i} E_{n-i,q^{w_2}}^{(h,r)}(w_1 x) T_{n,i,q^{w_1}}^{(h,r)}(w_2).
\end{aligned} \tag{2.9}
$$

Therefore, by (2.7) and (2.8), we obtain the following theorem.

Theorem 2.4. *For $n \geq 0$, and $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod 2$ and $w_2 \equiv 1 \pmod 2$, we have*

$$
\begin{aligned}
&\sum_{i=0}^n \binom{n}{i} [w_2]_q^i [w_1]_q^{n-i} E_{n-i,q^{w_1}}^{(h,r)}(w_2 x) T_{n,i,q^{w_2}}^{(h,r)}(w_1) \\
&= \sum_{i=0}^n \binom{n}{i} [w_1]_q^i [w_2]_q^{n-i} E_{n-i,q^{w_2}}^{(h,r)}(w_1 x) T_{n,i,q^{w_1}}^{(h,r)}(w_2),
\end{aligned}
$$

where

$$
T_{n,i,q}^{(h,r)}(w) = \sum_{j_1,\cdots,j_r=0}^{w-1} (-1)^{\sum_{l=1}^r j_l} q^{\sum_{l=1}^r (n+h-l-i)j_l} [j_1+\cdots+j_r]_q^i.
$$

References

[1] G. Andrews, B. C. Berndt, *Ramanujan's lost notebook, Part IV,* Springer, New York, 2013, xviii+439 pp.

[2] S. Araci, M. Acikgoz, *A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials*, Adv. Stud. Contemp. Math., 22(2012), no.3, 399-406.

[3] I. N. Cangul, V. Kurt, H. Ozden, Y. Simsek, *On the higher-order w-q-Genocchi numbers*, Adv. Stud. Contemp. Math. 19(2009), no. 1, 39-57.

[4] D. S. Kim, N. Lee, J. Na, K. H. Park, *Identities of symmetry for higher-order Euler polynomials in three variables (I)*, Adv. Stud. Contemp. Math. 22(2012), no. 1, 51-74.

[5] T. Kim, *Barnes-type multiple q-zeta functions and q-Euler polynomials*, J. Phys. A 43(2010), no. 25, 255201, 11pp.

[6] T. Kim, *Symmetry p-adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials*, J. Difference Equ. Appl. 14(2008), no.12, 1267-1277.

[7] T. Kim, *Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys. 16(2009), no. 1, 93-96.

[8] T. Kim, *An identity of the symmetry for the Frobenius-Euelr polynomials associated with the fermionic p-adic invariant q-integrals on $\mathbb{Z}_p$*, Rocky Mountain J. Math. 41(2011), no. 1, 239-247.

[9] Y.-H. Kim, K.-W. Hwang, *Symmetry of power sum and twisted Bernoulli polynomials*, Adv. Stud. Contemp. Math. 18(2009), no.2, 127-133.

[10] H. Ozden, Y. Simsek, S.-H. Rim, I. N. Cangul, *A note on p-adic q-Euler measure*, Adv. Stud. Contemp. Math. 14(2007), no.2, 233-239.

[11] S. H. Rim, J. Jeong, *On the modified q-Euler numbers of higher order with weight*, Adv. Stud. Contemp. Math. 22(2012), 93-98.

[12] Y. Simsek, *Interpolation functions of the Eulerian type polynomials and numbers*, Adv. Stud. Contemp. Math., 23(2013), no.2, 301-307.
[13] Y. Simsek, *Identities assoiciated with generalized Stirling type numbers and Eulerian type polynomials*, Math. Comput. Appl. 18(2013), no. 3, 251-263.
[14] H. J. H. Tuenter, *A symmetry of power sum polynomials and Bernoulli numbers*, Amer. Math. Monthly 108(2001), no. 3, 258-261.

Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea.
E-mail address: dskim@sogang.ac.kr

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea
E-mail address: tgkim2013@hotmail.com